

**Table 1. Evaluation of Lucento fungicide programs for control of early leaf spot in Oklahoma in 2023**

<b>Treatment<sup>1</sup>: Product, rate, timing<sup>2</sup></b>	<b>Early leaf spot (%) 30 Sep<sup>3</sup></b>	<b>Defoliation (%) 30 Sep<sup>4</sup></b>	<b>Yield (lbs./A)<sup>5</sup></b>
Untreated check	35.00 a	31.20 a	4,959 d
Lucento 5.5 fl. oz. (2, 4) Bravo 24 fl. oz. + Folicur 7.2 fl. oz. (3)	3.75 bc	3.25 c	5,612 a
Adastrio 8 fl. oz. (2, 4) Bravo 24 fl. oz. + Folicur 7.2 fl. oz. (3)	9.00 b	11.00 b	5,372 abc
Lucento 5.5 fl. oz. (2, 4) Headline 12 fl. oz. (3)	3.00 bc	3.50 c	5,300 abcd
Lucento 5.5 fl. oz. (2, 4) Abound 18.5 fl. oz.	2.50 bc	2.75 c	5,053 cd
Bravo 24 fl. oz. + Folicur 7.2 fl. oz. (1, 3) Lucento 5.5 fl. oz. (2, 4)	1.50 c	1.37 c	5,467 ab
Bravo 24 fl. oz. + Folicur 7.2 fl. oz. (1) Lucento 5.5 fl. oz. (2, 4) Headline 12 fl. oz. (3)	1.50 c	1.75 c	5,351 abcd
Bravo 24 fl. oz. + Folicur 7.2 fl. oz. (1) Lucento 5.5 fl. oz. (2, 4) Abound 18.5 fl. oz. (3)	1.75 c	2.00 c	5,460 ab
Bravo 24 fl. oz. + Folicur 7.2 fl. oz. (2) Abound 18.5 fl. oz. (3, 4)	8.25 bc	4.75 bc	5,184 bcd
LSD (P = 0.05) <sup>6</sup>	6.97	6.25	398.47
<sup>1</sup> Treatments 1 - 9 were applied with Induce @ 0.25% v/v. <sup>2</sup> Timing corresponds to the spray dates of 1=13 Jul, 2=31 Jul, 3=16 Aug, 4=1 Sep. <sup>3,4,5</sup> Values in a column followed by the same letter are not significant according to Fisher's least significant difference test at P=0.05. <sup>6</sup> Least significant difference.			