

Table 2. Efficacy of Priaxor and Provysol fungicides to manage early leaf spot in peanuts in 2023

Treatment ¹ : Product, rate, timing ²	Early leaf spot (%) 30 Sep ³	Defoliation (%) 30 Sep ⁴	Yield (lbs./A) ⁵
Untreated check	35.0 a	31.2 a	4,959
Bravo 24 fl. oz. (1, 5) Priaxor 8 fl. oz. (2, 4) Bravo 24 fl. oz. + Folicur 7.2 fl. oz. (3)	4.0 b	3.2 b	5,365
ABravo 24 fl. oz. (1, 5) Priaxor 8 fl. oz. (2, 4) Folicur 7.2 fl. oz. + Provysol 3 fl. oz. (3)	2.7 b	3.7 b	5,416
Bravo 24 fl. oz. (1, 5) Priaxor 8 fl. oz. (2, 4) Folicur 7.2 fl. oz. + Provysol 5 fl. oz. (3)	3.7 b	2.7 b	5,249
LSD (P = 0.05) ⁶	8.0	7.9	ns
¹ Treatments 1 – 5 were applied with Induce @ 0.25% v/v. ² Timing corresponds to the spray dates of 1 = 13 Jul, 2–31 Jul, 3 = 16 Aug, 4 = 1 Sep, 5 = 18 Sep. ^{3,4,5} Values in a column followed by the same letter are not significant according to Fisher's least significant difference test at P = 0.05. ⁶ Least significant difference.			