



Table 2. Peanut leaf spot and yield with Miravis® and Excalia® based fungicide programs at Fort Cobb, 2021^a.

Treatment	Application timing	102	117	139	153	153	Yield lb/A
		DAP	DAP	DAP	DAP	DAP	
		Florida Leaf Spot Scale					% Defol
Untreated		2.5	3.0	5.0	8.5	75	2590
Bravo®, Bravo®, Bravo®, Bravo®	P1, P2, P3, P4	1.0	2.0	3.8	5.3	26	3610
Bravo®, Miravis®, Bravo®, Miravis®	P1, P2, P3, P4	1.0	1.5	3.0	3.0	8	3640
Bravo®, Bravo®, Excalia®, Excalia®	P1, P2, P3, P4	1.5	2.5	4.8	6.3	58	3620
Bravo®, Bravo®, Excalia®, Bravo®	P1, P2, P3, P4	1.3	2.0	4.0	5.3	30	3260
Bravo®, Bravo®, Excalia®/Microthiol®, Bravo®	P1, P2, P3, P4	1.0	1.8	3.5	5.0	23	3430
LSD P=.10		0.5	0.7	0.5	1.2	9	520
CV (%)		30	25	10	17	20	13
Treatment Prob (F)		0.0008	0.0150	0.0001	0.0001	0.0001	0.0219

^aTreatment rates in fl oz/A: Bravo® Weather Stick (24.0), Miravis® (3.4), Excalia® (3.0); LSD = least significant difference, CV = coefficient of variation, DAP = days after planting, Defol = defoliation; Application Timings: POST1 (P1) = 60 DAP, POST2 (P2) = 76 DAP, POST3 (P3) = 91 DAP, POST4 (P4) = 102 DAP