



**Table 6. Gramoxone® rates, timings, and combinations effects on peanut stand at Fort Cobb, 2021.**

Treatments <sup>a</sup>	Rate (fl oz/A)	Timing	% Stand Reduction				
			33 DAP	47 DAP	61 DAP	77 DAP	92 DAP
Gramoxone®	10.8	14 DAC	1	1	1	1	0
+ Dual Magnum®	21.3	14 DAC	3	4	4	3	0
Gramoxone®	10.8	28 DAC	-	-	4	4	0
+ Dual Magnum®	21.3	28 DAC	-	-	1	1	0
Gramoxone®	21.6	14 DAC	1	1	1	1	0
+ Dual Magnum®	42.6	14 DAC	1	1	1	1	0
Gramoxone®	21.6	28 DAC	-	-	3	3	0
+ Dual Magnum®	42.6	28 DAC	-	-	3	3	0
Gramoxone®	10.8	14 + 28 DAC	1	1	1	1	0
+ Dual Magnum®	21.3	14 + 28 DAC	1	1	1	1	0
Gramoxone®	21.6	14 + 28 DAC	3	3	3	3	0
+ Dual Magnum®	42.6	14 + 28 DAC	4	4	4	4	0
GM® + DM® + Agri-Dex®	10.8 + 21.3 + 1%	28 DAC	-	-	3	1	0
GM® + DM® + Basagran®	10.8 + 21.3 + 16	28 DAC	-	-	3	3	0
LSD P=.10			NS	NS	NS	NS	NS
CV			148	135	125	135	0
Treatment Prob (F)			0.6311	0.4109	0.7112	0.8039	1.0000

<sup>a</sup>All treatments applied with Induce (0.25 %v/v) except where Agri-Dex® was applied. GM - Gramoxone®, DM = Dual Magnum®, DAC = days after cracking, LSD = least significant difference, CV = coefficient of variation, DAP = days after planting, NS = not significant