

OKLAHOMA

Perkins, Agronomy Research Station, Payne County
Rain-fed, Sown September 2004

Entry	2005				Total	Total NN*
	5/2	6/15	9/1	10/14		
Tons Dry Matter/Acre						
OK 49	1.37	0.92	1.18	1.42	4.90	4.91
OK 200-3-S-L	1.24	0.87	1.15	1.47	4.73	4.73
Good As Gold II	1.26	0.87	1.06	1.47	4.67	4.72
OK 200-3-C-M	1.27	0.82	1.13	1.39	4.61	4.59
6420	1.20	0.86	1.11	1.43	4.59	4.59
OK 200-3-S-M	1.23	0.84	1.10	1.41	4.57	4.58
55H05	1.18	0.89	1.08	1.45	4.59	4.49
HybrilForce-400	1.11	0.84	1.06	1.33	4.34	4.48
OK 200-3-C-S	1.21	0.80	1.09	1.35	4.44	4.44
362 HY	1.10	0.77	1.04	1.35	4.27	4.32
6530	1.05	0.78	1.06	1.38	4.28	4.18
Artesian Sunrise	0.97	0.90	1.01	1.36	4.23	4.14
Mean	1.18	0.85	1.09	1.40	4.52	4.52
5% LSD	0.19	0.06	0.07	0.10ns	0.29	0.23
CV (%)	14.0	5.8	5.5	6.1	5.5	4.5

Design: Randomized Complete Block
 No. of Reps: 6
 Experiment: 421
 ns = not significant at p= 0.05

Plot Size: 1x5m planted
 Plot Size: 1x5m harvested

*Total NN = Means adjusted by nearest neighbor analysis.
 Variety means are LSMEANS derived from nearest neighbor statistical analysis;
 therefore, season or multiple-year totals are not the arithmetic sum of individual cuts or
 years, respectively.